

Class 11th | Applied Maths



Unit : 1

Indices



$$a^m \times a^n = a^{m+n}$$

$$a^m \div a^n = a^{m-n}$$

$$a^m \times b^m = (a \times b)^m$$

$$a^m \div b^m = (a \div b)^m$$

$$a^0 = 1$$

$$a^{-m} = \frac{1}{a^m}$$

$$\frac{1}{a^{-m}} = a^m$$

$$a^m = b^m \Rightarrow a = b, m \neq 0$$

$$a^m = a^n \Rightarrow m = n, a \neq 1$$

$$\rightarrow \sqrt{3} = (3)^{1/2}$$

$$\rightarrow \sqrt[3]{5} = (5)^{1/3}$$

$$\rightarrow \sqrt[5]{7^2} = (7^2)^{1/5} = 7^{2/5}$$

$$\sqrt[n]{a^m} = a^{m/n}$$

$$(a^m)^n = a^{mn}$$

$$x^4 \times x^3 = x^7$$

$$\frac{x^4}{x^3} = x^{4-3}$$

$$a^{-m} = \frac{1}{a^m}$$

Simplify the following:

(i) $(3x^4y^3)(18x^3y^5)$

(ii) $\frac{3x^4y^3}{18x^3y^5}$

(iii) $\left(-\frac{2x^2}{y^3}\right)^3$

(iv) $\sqrt[3]{27^{-2}}$

$$\rightarrow 3x^4y^3 \times 18x^3y^5$$

$$\Rightarrow 54x^7y^8$$

$$\frac{3}{18} x^{4-3} y^{3-5}$$

$$\frac{x^1 y^{-2}}{6}$$

$$\frac{x}{6y^2}$$

$$\frac{(-2x^2)^3}{(y^3)^3}$$

$$\frac{-8x^6}{y^9}$$

$$(27^{-2})^{1/3}$$

$$[(3^3)^{-2}]^{1/3}$$

$$[3^{3 \times -2}]^{1/3}$$

$$3^{3 \times (-2) \times \frac{1}{3}}$$

$$3^{-2} = \frac{1}{3^2} = \frac{1}{9}$$

$$(3)^{\frac{1}{12}}$$

$$\frac{3}{12} = \frac{1}{4}$$

$$12 \frac{3 \times 1}{4 \times 3} = 12$$

$$\frac{1}{3} \xrightarrow{\times 4} 4$$

$$2CM(4,3) = 12$$

$$(i) \left[\left\{ (625)^{-\frac{1}{2}} \right\}^{-\frac{1}{4}} \right]^2$$

$$\sqrt[4]{28} \div \sqrt[3]{7}$$

$$\left[\left\{ (5^4)^{-\frac{1}{2}} \right\}^{-\frac{1}{4}} \right]^2$$

$$5^{4 \times -\frac{1}{2} \times -\frac{1}{4} \times 2}$$

$$5^1 = 5$$

$$(28)^{\frac{1}{4}} \div (7)^{\frac{1}{3}}$$

$$\frac{28^{\frac{1}{4}}}{7^{\frac{1}{3}}}$$

$$\frac{(28)^{\frac{3}{12}}}{(7)^{\frac{4}{12}}}$$

$$\frac{(28^3)^{\frac{1}{12}}}{(7^4)^{\frac{1}{12}}}$$

$$\left(\frac{28^3}{7^4} \right)^{\frac{1}{12}}$$

$$\left(\frac{28 \times 28 \times 28}{7 \times 7 \times 7 \times 7} \right)^{\frac{1}{12}}$$

$$\left(\frac{4 \times 4 \times 4}{7} \right)^{\frac{1}{12}}$$

$$\left(\frac{64}{7} \right)^{\frac{1}{12}}$$

$$\frac{a}{b} \div \frac{c}{d} \Rightarrow \boxed{\frac{a}{b} \times \frac{d}{c}}$$

$$\frac{a}{b} \times \frac{c}{d} \times \frac{e}{f} \rightarrow \underline{\text{Asam}}$$

Simplify:

$$\left(\frac{81}{16}\right)^{-\frac{3}{4}} \times \left[\left(\frac{25}{9}\right)^{-\frac{3}{2}} \div \left(\frac{5}{2}\right)^{-3}\right]$$

$$\left(\frac{3^4}{2^4}\right)^{-\frac{3}{4}} \times \left[\left(\frac{5^2}{3^2}\right)^{-\frac{3}{2}} \div \left(\frac{5}{2}\right)^{-3}\right]$$

$$\frac{3^{4 \times -\frac{3}{4}}}{2^{4 \times -\frac{3}{4}}} \times \left[\frac{5^{2 \times -\frac{3}{2}}}{3^{2 \times -\frac{3}{2}}} \div \frac{5^{-3}}{2^{-3}}\right]$$

$$\left(\frac{3^{-3}}{2^{-3}}\right) \times \left[\frac{5^{-3}}{3^{-3}} \div \frac{5^{-3}}{2^{-3}}\right]$$

$$\frac{2^3}{3^3} \times \left[\frac{3^3}{5^3} \div \frac{2^3}{5^3}\right]$$

$$\frac{2^3}{3^3} \times \frac{3^3}{5^3} \times \frac{5^3}{2^3}$$

①

$$2^3 \times 3^3$$

$$\begin{array}{r} 2 \overline{) 216} \\ 2 \overline{) 108} \\ 2 \overline{) 54} \end{array}$$

$$\begin{array}{r} 3 \overline{) 27} \\ 3 \overline{) 9} \\ 3 \overline{) 3} \end{array}$$

$$36 \times 4 = 144$$

Find the value of

$$\frac{4}{(216)^{-\frac{2}{3}}} + \frac{1}{(256)^{-\frac{3}{4}}} + \frac{2}{(243)^{-\frac{1}{5}}}$$

$$\frac{4}{(2^3 \times 3^3)^{-\frac{2}{3}}} + \frac{1}{(2^8)^{-\frac{3}{4}}} + \frac{2}{(3^5)^{-\frac{1}{5}}}$$

$$\Rightarrow \frac{4}{(2^{\cancel{3}^{-2}} \times 3^{\cancel{3}^{-2}})} + \frac{1}{2^{\cancel{8}^{-3}}} + \frac{2}{3^{\cancel{5}^{-1}}}$$

$$\Rightarrow \frac{4}{2^{-2} \times 3^{-2}} + \frac{1}{2^{-6}} + \frac{2}{3^{-1}}$$

$$4 \times 2^2 \times 3^2 + 2^6 + 2 \times 3$$

$$\Rightarrow 4 \times 4 \times 9 + 64 + 6$$

$$144 + 64 + 6$$

$$\underline{214}$$

$$0.01 = \frac{1}{100} = \frac{1}{10^2}$$

Evaluate: $\sqrt{\frac{1}{4}} + (0.01)^{-\frac{1}{2}} - (27)^{\frac{2}{3}}$. Leave your answer as a fraction.

$$\left(\frac{1}{4}\right)^{\frac{1}{2}} + (0.01)^{-\frac{1}{2}} - (3^3)^{\frac{2}{3}}$$

$$\frac{1^{\frac{1}{2}}}{(2^2)^{\frac{1}{2}}} + \left(\frac{1}{10^2}\right)^{-\frac{1}{2}} - 3^{3 \times \frac{2}{3}}$$

$$\Rightarrow \frac{1}{2} + \frac{1}{(10^2)^{-\frac{1}{2}}} - 3^2$$

$$\frac{1}{2} + \frac{1}{10^{-1}} - 9$$

$$\frac{1}{2} + 10 - 9$$

$$\frac{1}{2} + 1$$

$$\frac{3}{2}$$

Simplify:

$$\left(\frac{1}{4}\right)^{-2} - 3(8)^{\frac{2}{3}}(4)^0 + \left(\frac{9}{16}\right)^{-\frac{1}{2}}$$

$$\frac{(1)^{-2}}{4^{-2}} - 3 \times (2^3)^{\frac{2}{3}} \times 4^0 + \left(\frac{3^2}{4^2}\right)^{-\frac{1}{2}}$$

$$\frac{1}{4^{-2}} - 3 \times 2^2 \times 1 + \frac{(3^2)^{-\frac{1}{2}}}{(4^2)^{-\frac{1}{2}}}$$

$$\Rightarrow 4^2 - 12 + \frac{3^{-1}}{4^{-1}}$$

$$\Rightarrow 16 - 12 + \frac{4}{3}$$

$$4 + \frac{4}{3}$$

$$\frac{16}{3}$$

$$2^{3+4} = 2^3 \times 2^4$$

$$2^{x+y} = 2^x \cdot 2^y \Rightarrow 2^{m+2} = 2^m \cdot 2^2$$

Simplify the following:

(i) $\frac{5^{n+2} - 6 \cdot 5^{n+1}}{13 \cdot 5^n - 2 \cdot 5^{n+1}}$

(ii) $\left(\frac{x^m}{x^n}\right)^{m+n} \left(\frac{x^m}{x^l}\right)^{n+l} \left(\frac{x^l}{x^m}\right)^{l+m}$

(iii) $\left(x^{\frac{1}{3}} + x^{-\frac{1}{3}}\right) \left(x^{\frac{2}{3}} - 1 + x^{-\frac{2}{3}}\right)$

$$\Rightarrow \frac{5^n \cdot 5^2 - 6 \cdot 5^n \cdot 5^1}{13 \cdot 5^n - 2 \cdot 5^n \cdot 5^1}$$

$$\Rightarrow \frac{5^n [5^2 - 6 \times 5]}{5^n [13 - 10]}$$

$$\Rightarrow \frac{25 - 30}{3} = -\frac{5}{3}$$

$$\begin{aligned} & (x^{m-n})^{m+n} \cdot x^{(n-1)(n+1)} \cdot x^{(1-m)(1+m)} \\ & x^{m^2-n^2} \times x^{n^2-1^2} \times x^{1^2-m^2} \\ & x^{\cancel{m^2} - \cancel{m^2} + \cancel{n^2} - \cancel{n^2} + \cancel{1^2} - \cancel{1^2} - \cancel{m^2} + \cancel{m^2}} \\ & x^0 \\ & 1 \end{aligned}$$

$$(x^{1/3} + x^{-1/3})(x^{2/3} + x^{-2/3} - 1)$$

$$x^{1/3} \times x^{2/3} + x^{1/3} \times x^{-2/3} - x^{1/3} + x^{-1/3} \times x^{2/3} + x^{-1/3} \times x^{-2/3} - x^{-1/3}$$

$$x^{1/3+2/3} + x^{1/3-2/3} - x^{1/3} + x^{-1/3+2/3} + x^{-1/3-2/3} - x^{-1/3}$$

$$x^{3/3} + \cancel{x^{-1/3}} - \cancel{x^{1/3}} + \cancel{x^{1/3}} + x^{-3/3} - \cancel{x^{-1/3}}$$

$$x + x^{-1}$$

$$x + \frac{1}{x}$$

If $a = b^{2x}$, $b = c^{2y}$ and $c = a^{2z}$, prove that $xyz = 1/8$.

$$a = b^{2x}$$

$$a = (c^{2y})^{2x}$$

$$a = c^{4xy}$$

$$a = (a^{2z})^{4xy}$$

$$\underline{a} = \underline{a}^{8xyz}$$

$$1 = 8xyz$$

$$\frac{1}{8} = xyz$$

$$2^x = 3^y = 4^z$$

$$2 \times \frac{1}{y}$$

If $a^x = b^y = c^z$ and $b^2 = ac$ prove that $y = \frac{2xz}{z+x}$.

$$\left(\frac{2}{y}\right) = \left(\frac{2+x}{2z}\right) \Rightarrow \frac{y}{2} \rightarrow \frac{2z}{2+x} \Rightarrow y = \frac{2xz}{2+x}$$

$$a^x = b^y = c^z = k$$

$$a^x = k$$

$$b^y = k$$

$$c^z = k$$

$$(a^x)^{\frac{1}{x}} = (k)^{\frac{1}{x}}$$

$$a = k^{\frac{1}{x}}$$

$$(b^y)^{\frac{1}{y}} = (k)^{\frac{1}{y}}$$

$$b = k^{\frac{1}{y}}$$

$$(c^z)^{\frac{1}{z}} = (k)^{\frac{1}{z}}$$

$$c = k^{\frac{1}{z}}$$

$$b^2 = ac$$

$$(k^{\frac{1}{y}})^2 = k^{\frac{1}{x}} \times k^{\frac{1}{z}}$$

$$k^{\frac{2}{y}} = k^{\frac{1}{x} + \frac{1}{z}}$$

$$\frac{2}{y} = \frac{1}{x} + \frac{1}{z}$$

$$\frac{a \cdot b}{1} = \frac{1}{c} \quad \frac{1}{ab} = c$$

$$\frac{1+b}{ab}$$

$$\frac{1}{ab} + \frac{1}{c}$$

If $abc = 1$, show that

$$\frac{1}{1+a+b^{-1}} + \frac{1}{1+b+c^{-1}} + \frac{1}{1+c+a^{-1}} = 1$$

$$\begin{aligned} & \frac{1}{1+a+\frac{1}{b}} + \frac{1}{1+b+\frac{1}{c}} + \frac{1}{1+\frac{1}{a}+\frac{1}{b}} \\ & \frac{1}{\frac{b+ab+1}{b}} + \frac{1}{1+b+\frac{1}{c}} + \frac{1}{\frac{1}{1}+\frac{1}{ab}+\frac{1}{a}} \\ & \frac{b}{1+ab+b} + \frac{1}{1+b+\frac{1}{c}} + \frac{ab+1+b}{ab} \\ & \frac{b}{1+ab+b} + \frac{1}{1+ab+b} + \frac{ab}{1+ab+b} \end{aligned}$$

$$\frac{\cancel{b+1+ab}}{\cancel{1+ab+b}} = 1$$

$$x^3 = [(28)^{\frac{1}{3}}]^3 = 28$$

$$x^3 - y^3 = (x - y)(x^2 + y^2 + xy)$$

If $x = \sqrt[3]{28}$ and $y = \sqrt[3]{27}$, find the value of $x + y - \frac{1}{x^2 + xy + y^2}$.

$$x = (28)^{\frac{1}{3}} \quad y = (27)^{\frac{1}{3}}$$

$$\Rightarrow (x + y) - \frac{(x - y)}{(x - y)(x^2 + xy + y^2)}$$

$$\Rightarrow (x + y) - \frac{(x - y)}{x^3 - y^3}$$

$$\begin{aligned} x^3 &= 28 \\ y^3 &= 27 \end{aligned}$$

$$(x + y) - \frac{(x - y)}{28 - 27}$$

$$(x + y) - (x - y)$$

$$x + y - x + y$$

$$\begin{aligned} 2y &= 2(27)^{\frac{1}{3}} \\ &= 2 \times (3^3)^{\frac{1}{3}} \\ &= 2 \times 3 = 6 \end{aligned}$$

Given $1176 = 2^p \cdot 3^q \cdot 7^r$ find

$$1176 = 2^p \cdot 3^q \cdot 7^r \quad p, q, r$$

$$\rightarrow 2 \times 2 \times 2 \times 3 \times 7 \times 7 = 2^p \times 3^q \times 7^r$$

$$2^3 \times 3^1 \times 7^2 = 2^p \times 3^q \times 7^r$$

$$p = 3$$

$$q = 1$$

$$r = 2$$

2	1176
2	588
2	294
3	147
7	49
7	7

$$\frac{1}{2} \times 27$$

$$4 \times 27 = \frac{108}{125}$$

Solve the following equations for x:

(i) $4^{2x} = \frac{1}{32}$

(ii) $\sqrt{\left(\frac{3}{5}\right)^{1-2x}} = 4\frac{17}{27}$

$$(2^2)^{2x} = \frac{1}{2^5}$$

$$2^{4x} = 2^{-5}$$

$$4x = -5$$

$$x = -\frac{5}{4}$$

$$\left[\left(\frac{3}{5}\right)^{1-2x}\right]^{\frac{1}{2}} = \frac{125}{27}$$

$$\left(\frac{3}{5}\right)^{\frac{1}{2}(1-2x)} = \frac{5^3}{3^3}$$

$$\left(\frac{3}{5}\right)^{\frac{1}{2}-x} = \frac{3^{-3}}{5^{-3}}$$

$$\left(\frac{3}{5}\right)^{\frac{1}{2}-x} = \left(\frac{3}{5}\right)^{-3}$$

$$\frac{1}{2} - x = -3$$

$$\frac{1}{2} + 3 = x$$

$$\frac{7}{2} = x$$

STAY CONNECTED

KEEP LEARNING

Thank You



Join our telegram channel
@NEXTTOPPERS_COMMERCE1

